Southwest Gastroenterology Associates 7788 Jefferson St NE, Albuquerque, NM 87109

Phone: (505) 999-1600 Fax: (505) 999-1650 southwestgi.com

MoviPrep Bowel Prep		
Date:	Time:	Physician:
Location: Lovelace Medical Center Downtown		
601 Dr. Martin Luther King Jr Ave NE, Albuquerque, NM 87102		
Please call to confirm appointment 505-999-1606 or 999-1610		

Follow the instructions below to make sure that your colon is completely clean before your colonoscopy. *You will need a driver on the day of your procedure*. Please call our office if you have any questions.

### **1 WEEK PRIOR TO YOUR PROCEDURE**

Let us know if you are on any of the following: Plavix, Coumadin, Effient, Eliquis, Pradaxa or Warfarin. Please call your managing provider for instructions on stopping your medication \_\_\_\_\_\_days prior to your procedure.

**Stop** taking iron supplements or vitamins with iron, fish oil, krill oil, shark oil supplements as well as fiber.

# **Shopping List**

- Pick up your prescription of MoviPrep bowel prep kit at your pharmacy
- Dulcolax (Bisacodyl) Laxative 5mg tablets. (4 tablets)
- Optional for a sore bottom plain or aloe baby wipes, Desitin, A&D ointment or Vaseline
- Clear liquids

### 3 days before your procedure

**No** fruits, vegetables, or corn. No foods that contain seeds like tomatoes, green chile, sesame seed burns, etc.

You can eat meat, fish, rice, pasta, bread without seeds, peanut butter, eggs, cheese, and cereal.

## 2 days before your procedure

**No** fruits, vegetables, or corn. **No** foods that contain seeds like tomatoes, green chile, sesame seed burns, etc. You can eat meat, fish, rice, pasta, bread without seeds, peanut butter, eggs, cheese, and cereal.

No solid foods after midnight

**Confirm** arrangements with your driver.

## 1 day before your procedure

Solid foods, milk, or milk product are **not** allowed. Drink only "clear liquids" for breakfast, lunch, and dinner. **PLEASE AVOID ANY RED OR PURPLE COLORED LIQUIDS.** Please read the labels.

### "CLEAR LIQUIDS" INCLUDE:

- Strained fruit juice without pulp (apple, white grape, lemonade)
- Water, tea, coffee (without milk or non-dairy creamer)
- Clear broth or chicken, beef or vegetable bouillon
- Plain Jell-O, Popsicles
- Alcohol is not permitted.

2:00 PM: take 4 Dulcolax/Bisacodyl tablets.

5:00 PM: begin drinking the first dose of MoviPrep solution. Drink 8 oz every 15 minutes.

9:00 PM: begin drinking the second dose of the prep solution. Drink 8 oz every 15 minutes.

### Day of procedure

**Take** you medications at least 2 hours prior to your procedure with a small sip of water only.

**Do not** have anything to eat or drink (no liquids) after midnight the night before your procedure.

**Do not** skip heart, blood pressure or seizure medications.

You will need to bring current insurance information card, picture ID, and copayment.